


A Directed Transformation of Stem Cells to Neurons by Furan-based Macrocyclic Small Molecules: Validation by RNA Transcriptomic Analysis

The Implications to Neurological Disorders

Javed Iqbal



August 16, 2025

What is a stem cell?

A single cell that can






Image prepared by Catherine Twomey for the National Academies, *Understanding Stem Cells: An Overview of the Science and Issues* from the National Academies, <http://www.nationalacademies.org/stemcells>. Academic noncommercial use is permitted.



Embryonic stem cells (ESCs) are pluripotent cells derived from the inner cell mass of a blastocyst, a very early stage embryo. They have the unique ability to differentiate into any cell type in the body (pluripotency) and to self-renew, making them valuable for research and potential therapeutic applications.

But there are ethical issues.....


Mesenchymal stem cells (MSCs) are multipotent cells that can differentiate into many types of specialized cells



*The process of differentiation can be influenced by a variety of substances, including: 4
Transcription factors, Transforming growth factor beta, and Wnt proteins*


Induced Pluripotent Cells (iPSC)

iPS Cells




Shinya Yamanaka, winner of the 2012 Nobel Prize in Medicine and Physiology (Yamanaka Factors)

Induced Pluripotent Stem Cells (iPSCs) are formed by 4 Transcription factors from somatic cells



Generation Of iPSC by Yamanaka Factors




Patient



Health disorder
treatment




Cell collection




Somatic Cells

Reprogramming



iPSCs

Genome editing



Genetically modified iPSCs

**iPSCs-derived research
and treatment**

Application



Differentiated cells

Differentiation



Disease modelling






Global
Stem Cell
Care

WHY STEM CELLS DIFFERENTIATION IMPORTANT


[/www.globalstemcellcare.com/](http://www.globalstemcellcare.com/)

iPSC





Shinya Yamanaka
Kyoto University

Yamanaka Transcription Factors






Chemical equivalent of a stem cell-a "pluripotent" organic functionality




Can Small molecules be used to produce iPS Cells?


Chemical approaches to studying stem cell biology!



Small Molecules versus Transcription factors mediated differentiation of Stem Cells






Small-molecule-mediated reprogramming: a silver lining for regenerative medicine




Synthetic small molecules that control stem cell fate

Some Examples of Small Molecules Directed Differentiation of Stem Cells

Structures of small molecules that promote reprogramming to generate iPSCs



Cells can be reprogrammed to iPSCs using viral or non-viral introduction of exogenous TFs or protein transfection




A small molecule accelerates neuronal differentiation in the adult rat

Heiko Wurdak, Sho utian Zhu, Kyung Hoon Min and Peter G. Schultz


September 7, 2010

107 (38) 16542-16547


<https://doi.org/10.1073/pnas.1010300107>



Our Foray into Small molecules induced trans-differentiation of Stem cell to Neuron like cells



Natural products as a source of inspiration



Our Chemical Toolbox-based Working Model



Key features in our design:

- 3D architectures
- sufficient complexity
- stereochemical and skeletal diversity
- synthesis in a reasonable time-scale
- easy to follow-up medicinal chemistry studies

- Cyclic compounds
- Large surface area
- Pre-organization
- Less freely rotating bonds
- Enhanced cell permeation


From Arya Research Team:

Chem. & Biol. 163 (2005)

Curr. Opin. Chem. Biol. 247² (2005)


Chem. Rev. 1999 (2009)

Introduction



Eribulin


- Anticancer drug approved by FDA (November 15, 2010) and marketed by Eisai
- Synthetic analog of a marine sponge natural product, *halichondrin B*
- Mechanistically, a unique inhibitor of microtubule dynamics




Halichondrin B

Three Key Fragments of Eribulin



A



B



C








ER-076349

From Kishi's Lab:
Bioorg. Med. Chem. Lett. **2004**, *14*, 5551-5554

Our Next Generation Approach (Oxy-Michael)

Sharpless
dihydroxylation






D-tartaric acid

Saidulu Konda








Goal: Develop a Practical and Scalable Method


Our Synthetic Path




Completion of Fragment A Synthesis



Our Second Approach (Iodocyclization)




Our Synthetic Path




1. 2N HCl, H₂O.THF
65 °C, 16 h, 85%

2. I₂, CH₃CN, -20 °C
3 h, 70%



Separable



+




Our Synthesis Path (EWG = -COOEt)



*mixture of diastereomers
(inseparable, 13:1)*

Our Key Iodocyclization



1. Ag_2O , DCBCl, DMF

TBAI, rt, 16 h, **70%**

2. LiAlH_4 , THF, 0 °C-rt




16 h, **75%**

3. TBDPSCI, Im, DCM


0 °C-rt, 3 h, **85%**

4. NaH, MeI, DMF, 0 °C

1 h, **80%**




Single diastereomer



Key Features in Our Building Macrocyclic Designs


1. Natural product-inspired 3D-architectures
2. Stereochemical and skeletal diversity
3. Practical and scalable synthesis
4. Easy structural modifications for undertaking medicinal chemistry projects

Eribulin Fragment-based Macrocyclic Diversity





Goal: Develop Modular Approaches to Obtain Different Macrocycles to Explore their Biological Properties

Synthesis of Key Fragments





1. vinylmagnesium bromide
CuI, HMPA, THF, -30 °C
3 h, 75%

2. Ag₂O, MeI, DMF, rt, 12 h
70%




1. vinylmagnesium bromide
CuI, HMPA, THF, -30 °C
3 h, 75%





2. Ag₂O, MeI, DMF, rt, 12 h
70%







1. vinylmagnesium bromide
CuI, HMPA, THF, -30 °C
3 h, 85%






Our Synthetic Path to Building A Macrocyclic Diversity (Approach 1)







1,3-trans Relationship-based Four Macrocycles




Diastereomeric Macrocyclic Diversity (Approach 2)









1,3-*cis* Relationship-based Four Macrocycles




Another Approach to Building the Macrocyclic Diversity







Derivatives Synthesized










Macrocyclic Diversity using 1,3-*Cis* (C29-C32) Tetrahydrofuran



Derivatives




Our Macrocyclic Synthesis Roadmap






Enantiomerically pure

Part 2: Stereoselective Synthesis of C14-C21 Fragment of Eribulin & Derived Macrocyclic Diversity

Naveen Kumar








Literature (Kishi Approach)




J. Am. Chem. Soc., **1992**, *114*, 3163



Our Synthetic Planning




L-(+)- Tartaric acid




Stereoselective Synthesis of Furan Ring




i. 2,2 DMP, PTSA
 MeOH, 60 °C, 16 h
 ii. LAH, THF, 16 h
60% 3 steps





i. NaH, BnBr, THF, 5 h
 ii. PPh₃, Im, I₂, THF, 6 h
70% 2 steps



i. CuI, vinylMgBr
 HMPA, -30 °C, 2 h
 ii. PTSA, THF, H₂O
 16 h, **58% 2 steps**





I₂, NaHCO₃
 Et₂O, H₂O, 4 h, **82%**



4:1 Separable diastereomers


nOe Studies



Major isomer

5

No nOe




Minor isomer

5'



nOe

Our Plan to Building the Macrocyclic Diversity




Natural Products with Tetrahydrofuran Ring

Macrocycle A







Haterumalides

	R_1	R_2	R_3	R_4
Haterumalide NA	Ac	H	OH	H
Haterumalide NB	Ac	H	O-n-Bu	H
Haterumalide NC	Ac	OH	O-n-Bu	H
Haterumalide ND	Ac	OH	OH	H
Haterumalide NE	H	H	OH	H
Haterumalide B	Ac	H		H

- Uemura and coworkers isolated haterumalides in 1999 from the Okinawan sponge *Ircinia* sp.
- Haterumalide NA exhibited cytotoxicity against P388 cells with an IC50 of 0.32 $\mu\text{g/mL}$ and moderate acute toxicity against mice with an LD99 of 0.24 g/kg.


Our Approach to Macrocycle A



Grubbs 2nd gen. cat.




17 membered macrocycles

Synthesis of Different Macrocycles Achieved




17 membered macrocycles

Our Synthesis Execution for Macrocycle B






18 membered macrocycles

Synthesis of Different Macrocycles Achieved







18 membered macrocycles

Synthesis of Fully De-protected Macrocycles Achieved



Part 3: Another Eribulin Fragment




Mahender Khatravath




August 16, 2025

Macrocyclic Diversity Synthesis Plan





Precise



Isomeric A




Isomeric B





17 Membered Macrocycles

Our Synthetic Planning




Our Synthetic Path



L-Ascorbic acid



i. Acetone, CuSO_4 , rt, 24 h
ii. 30% H_2O_2 , K_2CO_3 , H_2O
rt, 24 h

iii. EtI , CH_3CN , 70°C , 24 h
80% 3 steps




i. Ag_2O , BnBr , Toluene
rt, 2 h, 90%

ii. LAH, THF, 12 h, rt, 90%
iii. $(\text{COCl})_2$, Et_3N , DMSO
DCM, -78°C , 1 h




DBU, NaI, THF, -78°C
2 h, 69% 2 steps




i. LiOH, THF-MeOH
rt, 1 h

ii. TFA- H_2O (9:1), 0°C -rt
7 h, 72% 2 steps





OsO_4 , Acetone


H_2O , rt, 2 h
82%



Ac_2O , DMAP, Py
rt, 16 h, 48%





Deviation in Our Approach




A

TFA, H₂O
ACN, 70 °C
1 h, 70%





TBSCl, Im
DCM, 12 h
40%




B






TFA, H₂O
ACN, 70 °C
1 h, 70%




TBSCl, Im
DCM, 12 h
40%




Stereocontrolled Michael!





Macrocyclic Diversity- An Example




Macrocyclic Diversity- An Example



Neural Differentiation with Small Molecules



The regenerative repair of neurons in acute injuries and neurodegeneration diseases







Reprogramming cell fates by small molecules

Neuronal stem cells


Neurons

Death of neurons





Neurotrophic compounds


Synthetic small molecules that induce neuronal differentiation in neuroblastoma and fibroblast cells, *Mol. BioSyst.*, 2015,11, 2727-2737





Neuroblastoma cells



Neurogenic cells






Fibroblast cells





Neurogenic cells

Small molecule-based lineage switch of human adipose-derived stem cells (hADSC) into neural stem cells and functional GABAergic neurons (iGN, iN)


Scientific Reports volume 7, Article number: 10166 (2017)



NEUROGENESIS



Our Studies




Chemical Approaches to Trans-differentiation of Human Mesenchymal Stem Cells (from Healthy Donors) to Neurons ; Our Approach



Mesenchymal Stem Cells seeded onto a 60 mm plate (~3 million cells/well)



The cells are allowed to reach ~80-90% confluency after which corresponding media is added to each well



Medium in wells is changed every 72 hr









Towards the end of the induction, the cells are scraped and saved for further analysis and characterization


Cells grown in different growth media are used for RT-PCR analysis to determine the expression levels of lineage specific genes

Adopted from Transcell's IP:
PCT/IB2014/000226
WO/2014/ 135950 PCT




Small Molecule-directed Differentiation of Stem Cells to Neuron-like Cells



iPS cells are stem cells created from a patient's own cells, such as skin or blood.





Small molecule induced trans-differentiation of human stem cells to neuron-like cells




Hierarchical clustering analysis:

Selected neuro-development genes up-regulation:

With 17a triplicates







GAPDH mRNA is present in brain

NEFM is a biomarker of neuronal damage

RTN4 expressed during neuronal development by neurons




Next Generation Sequencing - RNA Transcriptomic Studies




Collaborative work:

- Arya Group
- Transcell Biologics Hyderabad
- MedGenome Labs Bangalore




Control Vs Furan, 17a	Up regulated	Down regulated
Control Vs Furan, 17a	1372	1325

Triplicate studies



Hierarchical clustering




Hierarchical Clustering is visually represented as Dendrogram:


It is a method for grouping objects. The objects within a group are like each other and different from objects in other groups.

Effect on Pro-and Anti-Apoptosis Genes


Triplicate studies



Apoptosis mediates the precise and programmed natural death of neurons and is a physiologically important process in neurogenesis during maturation of the CNS.




Effect on Neurogenesis and Neurodevelopment




Neurogenesis:
new neurons are formed in the brain

Up-regulated genes upon treatment with 17a




Neurodevelopment:
Brain's development of neurological pathways that influence performance or functioning.



August 16, 2025

Neurogenesis Biomarkers



GABA input
 Glutamate input
 Glutamate output

Pax6^(52,62,70)
 Sox2⁽⁶⁴⁻⁶⁹⁾
 Hes6⁽⁷²⁾
 REST⁽⁷⁵⁻⁷⁷⁾
 TAp73⁽¹³⁹⁻¹⁵⁵⁾
 p53¹³¹⁻¹³⁷

Proliferation
 Self-renewal

Pax6^{71,72}
 SOX3⁸⁴
 SOX11⁸⁵⁻⁸⁷
 Tbr2^(10, 64)
 FoxG1⁸⁸⁻⁸⁹
 NeuroD1⁹²
 TAp73¹³⁹⁻¹⁵⁵

Differentiation
 Survival
 Migration

Prox1⁸¹⁻⁸³
 CREB⁹⁰⁻⁹⁴
 Tbr1^(10, 64)
 Ascl1/Mash1^(10, 64)

Prox1⁸¹⁻⁸³
 REST⁷⁵⁻⁷⁷
 Tbr1^(10, 64)
 TAp73¹³⁹⁻¹⁵⁵



Terminal
 differentiation

CREB⁹⁰⁻⁹⁴

Synaptogenesis

Mechanism of Neurogenesis: WNT Pathway


The complex of WNT, LRP, and Frizzled promotes the translocation of β -catenin into the nucleus, resulting in the transcription of β -catenin target genes.




Transcriptome data clearly showed that 17a up-regulated genes related to neurogenesis, axaongogenesis, neurodifferentiation, neurodevelopment and neuronal migration.

- **As shown 17a up-regulated genes NTRK2, CNTN1, NDNF, RET, CAMK1D and ATP8A2 which are known to be involved in neuronal development and differentiation.**
- **An Increased PCSK9 expression is also known to inhibit the amyloid β peptide ($A\beta$)-induced neuronal apoptosis by reducing the $A\beta$ generation**
- **The genes involved in neurogenesis, such as, CHAC1, CHAD7, PAX6, PCSK9, PHGDH, PRDM6 were upregulated.**
- **Several critical genes (for example, BCL2, TGFB2 and PCSK9) that are responsible for anti-apoptosis were found to be up-regulated**


Neuron's role in Neurological disorders (Alzheimer's and Parkinson)



a
Healthy brain



b
Alzheimer's brain




DNA methylation landscape in induced pluripotent stem cells (iPSCs) and directly converted neurons (iNs).

OKSM pathway

Reprogramming of aged fibroblasts to the pluripotent cell (iPSCs) stage is characterized by methylation erasure and reset of epigenetic clock.

OKSM (OCT4, KLF4, SOX2, C-MYC);


BAM (ASCL1, BRN2, MYT1L).




BAM Pathway

iNs derived from aged fibroblasts bypass the intermediate pluripotent cell stage, but still lack a faithful CH DNA methylation and show an altered CpG hypermethylation pattern at promoter regions.

Our Studies on Small Molecule based differentiation of iPSC derived from samples of Alzheimer's and Parkinson patients



Up-regulation of Genes That are known to be Down Regulated in Patients with Alzheimer's





CHGB gene: Encodes a tyrosine-sulfated secretory protein abundant in neurons

Upon treatment with small molecule, 17a

Upregulation of genes in Alzheimer's patient's stem cells during chemical trans-differentiation

- **SCN3B Gene:** Responsible for generation and propagation of action potentials in neurons and muscle
- **CD200 Gene:** negative regulation of neuroinflammatory response and neuron death
- **Coro1A Gene:** negative regulation of neuron apoptotic process
- **EPB41L3 Gene:** neuron projection morphogenesis
- **BRSK2 Gene:** neuron differentiation

Up-regulation of Genes That are known to be Down Regulated in Patients with Parkinson's



SCN1A gene: Provides instructions for making one part of sodium channel NaV 1.1 transmitting signals in neurons found in the brain.

Upon treatment with small molecule, 17a


Upregulation of genes in Parkinson's patient's stem cells during chemical trans-differentiation

- **DDC Gene:** Help in producing dopamine and serotonin neurotransmitter
- **GABRA2 Gene:** receptor in mammalian brain
- **KCND3 Gene:** Voltage gated K channel regulating neurotransmitter release and neuronal excitability

In Conclusion.....

1. Natural products-inspired small molecules are good probes for trans- differentiation of iPSC to Neuron like cells.
2. Cells derived from Alzheimer's and Parkinson patients undergo small-molecules promoted differentiation of stem cells to neuron like cells.
3. These studies promise to be a viable Human iPSC-based therapy.


Role of Furan Macrocycles in modulating Wnt signalling pathway in Cancer stem Cell: A quick look




Activation of Wnt Signaling

Involvement of the Wnt- β -catenin-GSK3 signaling axis in the formation and maintenance of cancer stem cells (CSC) is responsible for tumor growth in several types of human malignancies

Inactive Wnt Signaling




Active Wnt Signaling



Wnt signaling regulates fate decisions in embryonic stem cells.


1. GSK3 β acts as a tumor suppressor by curbing canonical Wnt- β -catenin signaling.

2. The small molecule (BIO) inhibits GSK3 β and thus increases the activity of β -catenin.




Wnt signaling in cancer

Normal cells




Cancer cells




Genetic Aberrations

Mutation of tumor suppressor genes




Epigenetic modifications


DNA methylation




Histone modifications



Non-coding RNAs



Activation of Wnt/ β -catenin pathway




Agents/Phytochemicals




Wnt inhibiting molecules




Crocin




Curcumin



Acetyl shikonin




ACBP



Clinical applications




DK419



IWR-1




Wnt Luciferase Screening Data




a cell-based test that uses a luciferase gene to measure the activity of the Wnt/ β -catenin signaling pathway in cultured cells:

Five
Analog



SM-2015-61/62/63/64/65





SM-2015-66/67/68/69/70

Family 1 Hits

Effect on Tumorspheres (G2*) from Patient Buccal Mucosa (Indian Patient 1)

G2 = 2nd generation



In Summary:

The studies involving furan-based macrocyclic molecules clearly highlight the potential therapeutic utility for such inhibitors to mediate **Wnt–GSK3– β -catenin axis activity at the expense of Cancer Stem Cells self-renewal and survival.**

Sincere Thanks to...



The Arya Group
DRILS Hyderabad

Transcell Biologics
Hyderabad

MedGenome Labs
Bangalore



MoES/CSIR-CDRI
*Science and Technology Park
University of Pune*



Thank You